CASIMERSEN (AMONDYS 45 [®]) PRESCRIBER ORDER FORM					
Fax completed form, insurance information, and clinical documentation to: (410) 558-6439					
	Patient Name:			Date of Birth:	
option care health [,]	Address:				
	Phone:	Height: Weight: Ibs Inches cm Date weight obtained:			
Clinical Information					
Primary Diagnosis Description: Duchenne muscular dystrophy (DMD) ICD-10 Code: G71.01					
Allergies:					
Casimersen (AMONDYS 45 [®]) Prescription					
Casimersen (AMONDYS 45 [®]) refill as directed x 1 year					
Infuse 30 mg/kg IV over 35 to 60 minutes every week (+/- 3 days to allow for patient/nurse scheduling).					
Dose will be rounded to closest 100 mg.					
Flush IV tubing with NS 10 to 20 mL after each infusion.					
Prescriber will obtain weight for non-ambulatory patients and provide dose changes to pharmacy as needed. Prescriber will arrange monthly dipstick proteinuria monitoring.					
Ancillary Orders Anaphylaxis Kit					
If this is a 1 st infusion dose, would you like Option Care Health to provide an anaphylaxis kit with the 1 st dose?					
□ Yes □ No					
Dosage: • Epinephrine 0.3 mg (> 30 kg), 0.15 mg (15 to 30 kg), or 0.01 mg/kg (< 15 kg) SQ or IM x 1; repeat x 1 in 5 to 15 min PRN.					
 Diphenhydramine 25 mg (> 30 kg) or 1.25 mg/kg (≤ 30 kg) IV or IM; repeat x 1 in 15 min PRN no improvement 					
 Normal saline 500 mL (> 30 kg) or 250 mL (≤ 30 kg) IV at KVO rate PRN anaphylaxis. Patients ≤ 30 kg, infuse over 2 to 4 hours PRN headache rated > 5 on pain scale. 					
Medication Orders					
Lidocaine/prilocaine 2.5%/2.5% (or equivalent) anesthetic cream 30 gm – apply topically 30 min prior to venipuncture or port access as needed for numbing.					
□ Other:					
IV Flush Orders					
 Peripheral: NS 2 to 3 mL pre-/post-use. Implanted Port: NS 5 to 10 mL pre-/post-use and 10 to 20 mL pre-/post-lab draw. Heparin (100 unit/mL) 3 to 5 mL post-use. For maintenance, heparin (100 unit/mL) 3 to 5 mL every 24 hr if accessed or weekly to monthly if not accessed. If unable to obtain implanted port access, it is acceptable to establish a peripheral IV access and administer peripherally. 					
Lab Orders Serum cystatin C and random urine protein-to-creatinine ratio, prior to infusion, every 3 months. No labs ordered at this time. Other:					
Skilled nurse to assess and administer and/or teach self-administration, where appropriate, via access device as indicated above. Nurse will provide ongoing support as needed. Refill above ancillary orders as directed x 1 year.					
I certify that the use of the indicated treatment is medically necessary and I will be supervising the patient's treatment.					
Prescriber Signature: Date:					
Prescriber Information					
Prescriber Name:			Phone:		Fax:
Address:			NPI:		
City, State: Zip:			Office Contact:		
CONFIDENTIAL HEALTH INFORMATION : Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that do not require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state laws. IMPORTANT WARNING: This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidentialit, he disclosure of which is governed by applicable law. If the reader of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.					