EDARAVONE (RADICAVA®) PRESCRIBER ORDER FORM						
Fax completed form, insurance information, and clinical documentation to: (888) 822-5060						
	Patient Name:			Dat	Date of Birth:	
option care health	Address:			·		
0,000	Phone:		Height:	$\square$ inches $\square$ cm	Weight:	☐ lbs ☐ kg
Clinical Information  Primary Diagnosis Description: Amyotrophic lateral sclerosis (ALS)  ICD-10 Code: G12.21						
Edaravone (Radicava®) Prescription						
Edaravone (Radicava®) 30 mg/100 mLs bags refill as directed x 1 year						
Initial Cycle:						
Maintenance Cycles:   Infuse 60 mg IV over 60 minutes daily for 10 days within a 14-day period followed by a 14-day drug-free period. Repeat maintenance cycle every 28 days.						
Ancillary Orders						
Anaphylaxis Kit  If this is a 1 <sup>st</sup> infusion dose, would you like Option Care Health to provide an anaphylaxis kit with the 1 <sup>st</sup> dose?  □ Yes □ No						
Dosage: • Epinephrine 0.3 mg (> 30 kg), 0.15 mg (15 to 30 kg), or 0.01 mg/kg (< 15 kg) SQ or IM x 1; repeat x 1 in 5 to 15 min PRN.						
• Diphenhydramine 25 mg (> 30 kg) or 1.25 mg/kg (≤ 30 kg) IV or IM; repeat x 1 in 15 min PRN no improvement.						
<ul> <li>Normal saline 500 mL (&gt; 30 kg) or 250 mL (≤ 30 kg) IV at KVO rate PRN anaphylaxis. Patients ≤ 30 kg, infuse over 2 to 4 hours PRN headache rated &gt; 5 on pain scale.</li> </ul>						
<b>Medication Orders</b>						
<ul> <li>□ Lidocaine/prilocaine 2.5%/2.5% (or equivalent) anesthetic cream 30 gm – apply topically 30 min prior to venipuncture or port access as needed for numbing.</li> <li>□ Other:</li> </ul>						
IV Flush Orders						
☐ <u>Peripheral</u>	<u>:</u>	Heparin (10 un	mL pre-/post-use. 0 unit/mL) 1 to 3 mL post-use. enance, heparin (10 unit/mL) 1 to 3 mL every 24 hr.			
☐ <u>Peripheral</u>	-Midline:	Heparin (100 u	ore-/post-use, 5 mL pre-lab draw, and 10 mL post-lab draw. unit/mL) 3 mL post-use. nce, flush with heparin (100 unit/mL) 3 mL every 24 hr.			
☐ PICC and C	entral Tunneled/Non-Tunneled	Heparin ☐ (10	pre-/post-use, 5 mL pre-lab draw, and 10 mL post-lab draw. $\Box$ (10 unit/mL) 5 mL $\underline{or}$ $\Box$ (100 unit/mL) 3 mL post-use. tenance, flush with heparin (10 unit/mL) 5 mL or (100 unit/mL) 3 mL every 24			
☐ <u>Implanted</u>	<u>Port:</u>	Heparin (100 u	nL pre-/post-use and 10 to 20 mL pre-/post-lab draw. 0 unit/mL) 3 to 5 mL post-use. nance, heparin (100 unit/mL) 3 to 5 mL every 24 hr if accessed or weekly to not accessed.			
	s and administer and/or teach s ort as needed. Refill above and			e, via access device	as indicated abov	e. Nurse will
I certify that the use of the indicated treatment is medically necessary and I will be supervising the patient's treatment.						
Prescriber Signature:  Prescriber I			Date:			
Prescriber Name:			Phone:		Fax:	
Address:			NPI:			
City, State:		Zip:	Office Contact:			

CONFIDENTIAL HEALTH INFORMATION: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that do not require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state laws. IMPORTANT WARNING: This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.